arXiv:1603.05535v4 [gr-qc] 23 Nov 2016

Chaplygin-gas Solutions of $f(R)$ Gravity

Maye Elmardi[1,2], Amare Abebe[3] and Abiy Tekola[4]

[1] Astrophysics, Cosmology and Gravity Centre, University of Cape Town, Rondebosch, 7701, South Africa

[2] Department of Mathematics and Applied Mathematics, University of Cape Town, Rondebosch, 7701, South Africa

[3] Department of Physics, North-West University, Mafikeng 2735, South Africa

[4] Las Cumbres Observatory Global Telescope Network, Goleta, CA 93117, USA

Abstract.

We explore exact $f(R)$ gravity solutions that mimic Chaplygin-gas inspired ΛCDM cosmology. Starting with the original, generalized and modified Chaplygin gas equations of state, we reconstruct the forms of $f(R)$ Lagrangians. The resulting solutions are generally quadratic in the Ricci scalar, but have appropriate ΛCDM solutions in limiting cases. These solutions, given appropriate initial conditions, can be potential candidates for scalar field-driven early universe expansion (inflation) and dark energy-driven late-time cosmic acceleration.

Keywords: Cosmology; $f(R)$ gravity; Chaplygin gas; dark energy; dark matter.

PACS numbers: 04.50.Kd, 04.25.Nx

1. Introduction

Explaining the not-yet-so-fully-understood accelerated expansion phases of the early and current stages of the Universe as well as the flatness in the rotational curves of spiral galaxies have certainly become in the top of to-do lists in modern cosmology and astrophysics. As a result, current research into the nature of so-called dark energy and dark matter has led to reconsiderations of the gravitational physics (General Relativity, GR) and cosmic fluid (baryonic matter) models underlying the standard cosmological model.

Among a large pool of alternative theories put forward to explain early-universe inflation [1] and late-time cosmic acceleration [2–8] are $f(R)$ *theories* of gravity (see Refs. [9–14] for more detailed reviews). Such theories were first speculated on by Buchdahl [15] but recent interest has centered on their potential candidacy as possible infrared (IR) and ultraviolet (UV) completions of GR (see, e.g., [16–21]). More recently, these gravitational alternative theories have found cosmological applications in, *inter alia*, the dynamical study of homogeneous cosmological models [22–38] and the linear growth of large-scale structures [39–47].

For quite sometime now, the Chaplygin gas [48] has been considered as another alternative cosmological model of a Friedmann-Lemaître-Robertson-Walker (FLRW) universe with an exotic perfect-fluid equation of state [49–54]. The model provides a cosmic expansion history consistent with a universe that transitions from a decelerating matter-dominated phase to a late-time accelerated one [52,55,56]. As a result, there have been several efforts to come up with a unified description of both dark matter and dark energy as manifestations of a single cosmic fluid that is the Chaplygin gas [52, 57–59].

The main objective of this work is to come up with a unified picture of both alternative cosmological scenarios discussed above. We study models of $f(R)$ gravity which, when we impose the Chaplygin gas equations of state (EoS) to their effective pressure and energy density, produce viable exact solutions that reduce to the ΛCDM scenario in the approximate cosmological limits.

The rest of this paper is organised as follows: In Sec. (2) we give a brief overview of $f(R)$ gravitational models and their corresponding field equations, and describe the background cosmic evolution under consideration. We give a brief description of the Chaplygin gas model and its generalized and modified EoS in Sec. (3). By comparing these EoS with the EoS for $f(R)$ models under certain assumptions, we deduce the form of the corresponding $f(R)$ Lagrangians. Finally in Sec. (4) we discuss the results and give our conclusions.

Natural units ($\hbar = c = k_B = 8\pi G = 1$) will be used throughout this paper, and Latin indices run from 0 to 3. The symbols ∇, $\tilde{\nabla}$ and the overdot $\cdot$ represent the usual covariant derivative, the spatial covariant derivative, and differentiation with respect to cosmic time. We use the $(-+++)$ spacetime signature and the Riemann tensor is defined by

$$R^a_{bcd} = \Gamma^a_{bd,c} - \Gamma^a_{bc,d} + \Gamma^e_{bd}\Gamma^a_{ce} - \Gamma^f_{bc}\Gamma^a_{df} \ , \tag{1}$$

where the Γ^a_{bd} are the Christoffel symbols (i.e., symmetric in the lower indices) defined by

$$\Gamma^a_{bd} = \frac{1}{2} g^{ae} \left(g_{be,d} + g_{ed,b} - g_{bd,e} \right) \ . \tag{2}$$

The Ricci tensor is obtained by contracting the *first* and the *third* indices of the Riemann tensor:

$$R_{ab} = g^{cd} R_{cadb} \ . \tag{3}$$

Unless otherwise stated, primes $'$ etc are shorthands for derivatives with respect to the Ricci scalar

$$R = R^a{}_a \tag{4}$$

and f is used as a shorthand for $f(R)$.

2. $f(R)$ Models

The action for $f(R)$ gravity can be given as

$$\mathcal{A} = \int \mathrm{d}^4x \sqrt{-g} \left[\frac{1}{2} f(R) + \mathcal{L}_m \right] \ , \tag{5}$$

where $f(R)$ is a general differentiable function of the Ricci scalar and $\mathcal{L}_m$ corresponds to the matter Lagrangian. For models of gravitation with such action, applying the variational principle with respect to the metric g_{ab} results in the generalized field equations

$$G_{ab} = T^m_{ab} + T^R_{ab} \equiv T_{ab} \ , \tag{6}$$

where T_{ab} is the total energy momentum tensor (EMT) and is conserved by virtue of the Einstein tensor $G_{ab} \equiv R_{ab} - \frac{1}{2} R g_{ab}$ being a covariantly conserved quantity. Here T^m_{ab} is the EMT of standard matter and

$$T^R_{ab} = \frac{1}{2} g_{ab} \left(f - R f' \right) + \nabla_b \nabla_a f' - g_{ab} \nabla_c \nabla^c f' + \left(1 - f' \right) G_{ab} \tag{7}$$

is the curvature contribution to the EMT. Since T^m_{ab} is a conserved quantity, we see from Eq. (6) that T^R_{ab} should also be conserved. In this setting, the linearized curvature components of the thermodynamical quantities are given by

$$\mu_R \equiv T^R_{ab} u^a u^b = \frac{1}{2}(R f' - f) - \Theta f'' \dot{R} + \frac{1}{3}(1 - f')\Theta^2 + f'' \tilde{\nabla}^2 R \ , \tag{8}$$

$$\begin{aligned} p_R \equiv \tfrac{1}{3}(T^R_{ab} h^{ab}) = {} & \frac{1}{6} \left(3f - R f' - 2R \right) + f'' \ddot{R} + f''' \dot{R}^2 + \frac{2}{3} \Theta f'' \dot{R} \\ & + \frac{1}{9}(1 - f')\Theta^2 - \frac{2}{3} f'' \tilde{\nabla}^2 R \ , \end{aligned} \tag{9}$$

$$q^R_a \equiv -T^R_{bc} u^b h^c_a = \frac{1}{3} f'' \Theta \tilde{\nabla}_a R - f''' \dot{R} \tilde{\nabla}_a R - f'' \tilde{\nabla}_a \dot{R} - \frac{2}{3}(1 - f') \tilde{\nabla}_a \Theta \ , \tag{10}$$

$$\pi^R_{ab} \equiv T^R_{cd} h^c{}_{\langle a} h^d{}_{b \rangle} = f'' \tilde{\nabla}_{\langle a} \tilde{\nabla}_{b \rangle} R \ , \tag{11}$$

and the total cosmic medium is composed of standard matter and the *curvature fluid*, with the total thermodynamical quantities given by

$$\mu \equiv \mu_m + \mu_R \ , \quad p \equiv p_m + p_R \ , \quad q_a \equiv q_a^m + q_a^R \ , \quad \pi_{ab} \equiv \pi_{ab}^R \ . \tag{12}$$

The trace of the Einstein field equations (6) gives

$$3f''\ddot{R} + 3\dot{R}^2 f''' + 3\Theta\dot{R}f'' - 3f''\tilde{\nabla}^2 R - Rf' + 2f + 3p_m - \mu_m = 0 \ , \tag{13}$$

and plays an important role as a constraint relation between matter and curvature.

Analogous to the matter EoS $p_m = w_m\mu_m$, where w_m is the matter EoS parameter, we can define the EoS for the curvature fluid as $p_R = w_R\mu_R$.

For FLRW spacetimes, the Ricci scalar R is given by

$$R = 2\dot{\Theta} + \frac{4}{3}\Theta^2 \ , \tag{14}$$

where Θ is the cosmic expansion parameter related to the cosmological scale factor $a(t)$ and the Hubble parameter $H(t)$ via the equations

$$\Theta \equiv 3\frac{\dot{a}(t)}{a(t)} = 3H(t) \ . \tag{15}$$

Solving for $\dot{\Theta}$ one can rewrite the above equation as

$$\dot{\Theta} = \frac{R}{2}\left(1 - \frac{4}{3R}\Theta^2\right) \ . \tag{16}$$

If the Ricci scalar varies slowly, i.e., if R is almost constant, the solution of this ordinary differential equation (o.d.e) takes the form

$$\Theta = \frac{1}{2}\sqrt{3R}\tanh\left[\sqrt{\frac{R}{3}}(t - t_0)\right] \ , \tag{17}$$

for some constant of integration t_0 that can be taken to be the time at the commencement of the inflationary phase of expansion. Solving for the cosmological scale factor gives

$$a(t) = a_0\sqrt{\cosh\left[\sqrt{\frac{R}{3}}(t_0 - t)\right]} \ . \tag{18}$$

For simplicity we set $t_0 \simeq 0$. During steady-state exponential expansion in a de Sitter spacetime (such as during inflation or late-time evolution), the approximation $\dot{\Theta} \to 0$ results in

$$R = \frac{4}{3}\Theta^2 = \text{const} \ , a(t) = a_0 e^{\frac{1}{3}\Theta t} \ , \tag{19}$$

and the matter content evolves according to

$$\mu_m = \mu_0\left(\frac{a(t)}{a_0}\right)^{-3(1+w_m)} \ . \tag{20}$$

During such an exponentially expanding cosmic evolution phase, for an initial energy density μ_0, we see that the matter energy density decays exponentially:

$$\mu_m = \mu_0 e^{-(1+w_m)\Theta t} \ . \tag{21}$$

3. Chaplygin-gas Solutions

Chaplygin-gas fluid models are perfect-fluid models currently posing as candidates to unify dark energy and dark matter. These fluid models were originally studied [48] in the context of aerodynamics, but only recently did they see cosmological applications [49–51, 57, 59, 60]. Among the interesting features of these models is that in the FLRW framework, a smooth transition between an inflationary phase, the matter-dominated decelerating era, and then late-time accelerated de Sitter phase of cosmic expansion can be achieved [52, 55].

3.1. Original and Generalized Chaplygin Gases

In the original treatment, the negative pressure associated with the Chaplygin gas models is related to the (positive) energy density through the EoS

$$p = -\frac{A}{\mu^{\alpha}} \tag{22}$$

for positive constant A and $\alpha = 1$. But this was later generalized [55, 61] to include $0 \leq \alpha \leq 1$. One of the first cosmological interpretations of such a fluid model was given in [62] where for flat universes, Eq. (22) corresponds to a viscosity term that is inversely proportional to dust energy density. Ever since the discovery of cosmic acceleration, however, both the original and generalized Chaplygin gas models have been extensively studied as alternatives to dark energy and/or unified dark energy and dark matter models (see, e.g., [50, 52, 55, 58–60, 63, 64]).

Now if we consider the background curvature energy density and isotropic pressure terms defined in Eqs. (8) and (9) above, in the constant-curvature limiting case, we have

$$\mu_R = \frac{1}{4}\left[R(f' + 1) - 2f\right] = -p_R \, . \tag{23}$$

This equation of state, with an effective EoS parameter $w_R = -1$, provides the condition for an exponential (accelerated) expansion with a constant Hubble parameter. The energy density μ_R (with its negative pressure p_R) remains constant and can be interpreted as playing the role of the cosmological constant Λ.

Considering the *curvature fluid* as a manifestation of the Chaplygin gas with the EoS (22), we obtain

$$p_R = -\mu_R = -\frac{A}{\mu_R^{\alpha}} \, , \tag{24}$$

which, using Eq. (23), leads to the o.d.e

$$R\frac{f(R)}{dR} - 2f(R) + R = 4A^{\frac{1}{\alpha+1}} \, . \tag{25}$$

Solving this o.d.e yields

$$f(R) = R + C_1 R^2 - 2A^{\frac{1}{\alpha+1}} \tag{26}$$

for an arbitrary (integration) constant C_1. We note that the ΛCDM solution $f(R) = R - 2\Lambda$ is already a particular solution with $C_1 = 0$ and $A = \Lambda^{\alpha+1}$. In particular, if $\alpha = 0$, then $A = \Lambda$, from which, going back to Eq. (24), one concludes $\mu_R = \Lambda$.

If we include the linearized Laplacian term in Eqs. (8) and (9) and use the eigenvalue $-\frac{k^2}{a^2}$ of the covariantly defined Laplace-Beltrami operator $\tilde{\nabla}^2$ on (almost) FLRW spacetimes

$$\tilde{\nabla}^2 R = -\frac{k^2}{a^2} R \tag{27}$$

for a comoving wavenumber k, we obtain the second-order o.d.e

$$B^2 R f''(R) - R f'(R) + 2f(R) - R + 4A^{\frac{1}{\alpha+1}} = 0 \,, \tag{28}$$

where here we have defined

$$B^2 \equiv \frac{4(2+3\alpha)}{3(1+\alpha)} \frac{k^2}{a^2} \,. \tag{29}$$

The solution of Eq. (28) is given, for arbitrary constants C_2, C_3, by

$$f(R) = R + C_2 \left[R^2 - 2RB^2\right] + C_3 \left[(R^2 - 2RB^2) Ei\left(1, -\frac{R}{B^2}\right) + (R - B^2) B^2 e^{\frac{R}{B^2}}\right] - 2A^{\frac{1}{\alpha+1}} \,, \tag{30}$$

which should reduce to the quadratic solution (26) for negligible values of B^2, i.e., for small first-order contributions to the energy density and pressure terms.

3.2. Modified Chaplygin Gas

Over the years, several modifications to the original and *generalized* Chaplygin gas models have been studied. If one considers the *modified Chaplygin gas* (MCG) EoS [61, 65–71]

$$p_R = \gamma \mu_R - \frac{A}{\mu_R^\alpha} \,, \tag{31}$$

then the resulting $f(R)$ model generalizes to

$$f(R) = R + C_4 R^2 - 2\left(\frac{A}{\gamma+1}\right)^{\frac{1}{\alpha+1}} \,, \tag{32}$$

where C_4 is an arbitrary integration constant.

The ΛCDM solution is a limiting case of this generalized model when $C_4 = 0$ and $A = (\gamma+1)\Lambda^{\alpha+1}$. In particular, if $\alpha = 0 = \gamma$, then $A = \Lambda$.

Following similar arguments as in the preceding subsection, if we include the linearized Laplacian contributions to the energy density and pressure, we get Eq. (28) generalized to

$$B^2 R f''(R) - R f'(R) + 2f(R) - R + 4\left(\frac{A}{\gamma+1}\right)^{\frac{1}{\alpha+1}} = 0\,, \tag{33}$$

the solution of which can be given by

$$f(R) = R + C_5\left[R^2 - 2RB^2\right] + C_6\left[(R^2 - 2RB^2)Ei\left(1, -\frac{R}{B^2}\right) + (R - B^2)B^2 e^{\frac{R}{B^2}}\right] - 2\left(\frac{A}{\gamma+1}\right)^{\frac{1}{\alpha+1}}\,, \tag{34}$$

for an arbitrary integration constants C_5 and C_6. This solution obviously generalizes Solutions (26),(30) and (32) and should reduce to the quadratic solution (26) for vanishingly small B^2 values. In [72], it has been shown that any quadratic Lagrangian leading to an isotropic, homogeneous cosmological model takes the form

$$f(R) = R - 2\Lambda - \frac{1}{6}\beta R^2\,, \tag{35}$$

where β is an arbitrary, real constant. If we keep only the quadratic solution in (34), i.e., if we set $C_6 = 0$, the Lagrangian (35) corresponds to the choice

$$C_5 = -\frac{1}{6}\beta\,, B = 0\,, A = (\gamma+1)\Lambda^{\alpha+1}\,. \tag{36}$$

Another interesting fact worth pointing out here is that the condition for the existence of a maximally symmetric vacuum solution in $f(R)$ gravity [72]

$$R_0 f'(R_0) = 2f(R_0) \tag{37}$$

leads to the quadratic solution resulting in the constraint

$$R_0\left(1 - 2C_5 B^2\right) - 4\left(\frac{A}{\gamma+1}\right)^{\frac{1}{\alpha+1}} = 0\,. \tag{38}$$

The corresponding GR de Sitter, anti-de Sitter and Minkowski solutions $R_0 = 4\Lambda$ (respectively for $\Lambda > 0$, $\Lambda < 0$ and $\Lambda = 0$) are obtained when $C_5 B^2 = 0$ and $A = (\gamma+1)\Lambda^{\alpha+1}$.

3.3. Modified Generalized Chaplygin Gas

The so-called *modified generalized Chaplygin gas* (mGCG) model is described by a barotropic equation of state of the form [59, 60]

$$p = \beta\mu - (1+\beta)\frac{A}{\mu^\alpha}\,. \tag{39}$$

Models of $f(R)$ gravity that satisfy the condition (23), at the sam time mimicking the mGCG, can be shown to be governed by the same equation as (25) and admit the same solutions (26), provided $\beta \neq -1$. On the other hand, if linearized Laplacian terms are included, then the corresponding differential equation in $f(R)$ generalizes to

$$D^2 R f''(R) - R f'(R) + 2f(R) - R + 4A^{\frac{1}{\alpha+1}} = 0 \,, \tag{40}$$

where we have defined

$$D^2 \equiv \frac{4\left[2 + 3\alpha + 3\beta(1+\alpha)\right]}{3(1+\alpha)(1+\beta)} \frac{k^2}{a^2} \,. \tag{41}$$

Worthy of note is that this equation and its solution

$$f(R) = R + C_7\left[R^2 - 2RD^2\right] + C_8\left[(R^2 - 2RD^2)Ei\left(1, -\frac{R}{D^2}\right) + (R - D^2)D^2 e^{\frac{R}{D^2}}\right] - 2\left(\frac{A}{\gamma+1}\right)^{\frac{1}{\alpha+1}} \,, \tag{42}$$

reduce to their *generalized* counterparts of Eqs. (28) and (30) when $\beta = 0$.

4. Discussions and Conclusion

In this paper we have explored models of $f(R)$ gravity that have Chaplygin gas equations of state. We started with the assumption that the effective energy density and isotropic pressure terms of Eqs. (8) and (9), sourced by the energy-momentum tensor of the curvature fluid (7), satisfy the equation of state of the Chaplygin gas, both in its original (Eq. 22) and generalized (Eq. 31) forms. We then reconstructed, for time-independent Ricci scalar, the f(R) Lagrangians that allow such equations of state. When only background (zeroth-order) terms are considered for the energy density and pressure, we notice that the effective equation of state is $w_R = -1$, and hence we naturally get $f(R)$ models that are quadratic in R but have limiting ΛCDM solutions.

On the other hand, keeping the first-order (Laplacian of R) terms in the energy density and pressure of the curvature fluid results in the effective energy density picking up linearized corrections to -1. The resulting linearized second-order o.d.es in $f(R)$ result in more generalized solutions including exponential integral terms, of which the quadratic Lagrangians are but a subset of solutions. These solutions also include ΛCDM as a limiting solution.

Acknowledgments

The authors thank John Barrow and Behnam Pourhassan for their constructive comments.

References

[1] Starobinsky, A. A. A new type of isotropic cosmological models without singularity. *Physics Letters B* **91**, 99–102 (1980).

[2] Carroll, S., Duvvuri, V., Turner, M. & Trodden, M. Is cosmic speed-up due to new gravitational physics? *Physical Review D* **70**, 043528 (2004).

[3] Magnano, G., Ferraris, M. & Francaviglia, M. Nonlinear gravitational lagrangians. *General relativity and gravitation* **19**, 465–479 (1987).

[4] Nojiri, S. & Odintsov, S. D. Modified gravity with negative and positive powers of curvature: Unification of inflation and cosmic acceleration. *Physical Review D* **68**, 123512 (2003).

[5] Sotiriou, T. P. & Liberati, S. Metric-affine $f(R)$ theories of gravity. *Annals of Physics* **322**, 935–966 (2007).

[6] Srivastava, S. Curvature inspired cosmological scenario. *International Journal of Theoretical Physics* **47**, 1966–1978 (2008).

[7] Capozziello, S., Cardone, V. & Troisi, A. Dark energy and dark matter as curvature effects? *Journal of Cosmology and Astroparticle Physics* **2006**, 001 (2006).

[8] Nojiri, S. & Odintsov, S. D. Modified gravity as an alternative for ΛCDM cosmology. *Journal of Physics A: Mathematical and Theoretical* **40**, 6725 (2007).

[9] De Felice, A. & Tsujikawa, S. $f(R)$ theories. *Living Rev. Rel* **13**, 1002–4928 (2010).

[10] Clifton, T., Ferreira, P. G., Padilla, A. & Skordis, C. Modified gravity and cosmology. *Physics Reports* **513**, 1–189 (2012).

[11] Capozziello, S. & De Laurentis, M. Extended theories of gravity. *Physics Reports* **509**, 167–321 (2011).

[12] Sotiriou, T. P. & Faraoni, V. $f(R)$ theories of gravity. *Reviews of Modern Physics* **82**, 451 (2010).

[13] Faraoni, V. $f(R)$ gravity: successes and challenges. *arXiv preprint arXiv:0810.2602* (2008).

[14] Bamba, K., Nojiri, S., Odintsov, S. D. & Sáez-Gómez, D. Inflationary universe from perfect fluid and $F(R)$ gravity and its comparison with observational data. *Physical Review D* **90**, 124061 (2014).

[15] Buchdahl, H. A. Non-linear Lagrangians and cosmological theory. *Monthly Notices of the Royal Astronomical Society* **150**, 1–8 (1970).

[16] Modesto, L. Super-renormalizable quantum gravity. *Phys. Rev. D* **86**, 044005 (2012).

[17] Modesto, L. & Rachwal, L. Super-renormalizable and finite gravitational theories. *Nuclear Physics B* **889**, 228–248 (2014).

[18] Biswas, T., Gerwick, E., Koivisto, T. & Mazumdar, A. Towards singularity and ghost free theories of gravity. *Phys.Rev.Lett.* **108**, 031101.

[19] Nojiri, S. & Odintsov, S. D. Introduction to modified gravity and gravitational alternative for dark energy. *International Journal of Geometric Methods in Modern Physics* **4**, 115–145 (2007).

[20] Nojiri, S. & Odintsov, S. D. Unified cosmic history in modified gravity: from $F(R)$ theory to lorentz non-invariant models. *Physics Reports* **505**, 59–144 (2011).

[21] Nojiri, S., Odintsov, S. & Sáez-Gómez, D. Cyclic, ekpyrotic and little rip universe in modified gravity. In *TOWARDS NEW PARADIGMS: PROCEEDING OF THE SPANISH RELATIVITY MEETING 2011*, vol. 1458, 207–221 (AIP Publishing, 2012).

[22] Rippl, S., van Elst, H., Tavakol, R. & Taylor, D. Kinematics and dynamics of $f(R)$ theories of gravity. *General Relativity and Gravitation* **28**, 193–205 (1996).

[23] Nojiri, S. & Odintsov, S. D. Modified $f(R)$ gravity consistent with realistic cosmology: From a matter dominated epoch to a dark energy universe. *Physical Review D* **74**, 086005 (2006).

[24] Bamba, K., Odintsov, S. D. *et al.* The future of the universe in modified gravitational theories: approaching a finite-time future singularity. *Journal of Cosmology and Astroparticle Physics* **2008**, 045 (2008).

[25] Bamba, K., Geng, C.-Q., Nojiri, S. & Odintsov, S. D. Crossing of the phantom divide in modified gravity. *Physical Review D* **79**, 083014 (2009).

[26] Amendola, L., Polarski, D. & Tsujikawa, S. Are $f(R)$ dark energy models cosmologically viable? *Physical review letters* **98**, 131302 (2007).

[27] Nojiri, S. & Odintsov, S. D. Modified gravity and its reconstruction from the universe expansion history. In *Journal of Physics: Conference Series*, vol. 66, 012005 (IOP Publishing, 2007).

[28] Carloni, S., Dunsby, P. K., Capozziello, S. & Troisi, A. Cosmological dynamics of R^n gravity. *Classical and Quantum Gravity* **22**, 4839 (2005).

[29] Capozziello, S., Nojiri, S., Odintsov, S. & Troisi, A. Cosmological viability of $f(R)$-gravity as an ideal fluid and its compatibility with a matter dominated phase. *Physics Letters B* **639**, 135–143 (2006).

[30] Carloni, S., Troisi, A. & Dunsby, P. Some remarks on the dynamical systems approach to fourth order gravity. *General Relativity and Gravitation* **41**, 1757–1776 (2009).

[31] Leach, J. A., Carloni, S. & Dunsby, P. K. Shear dynamics in Bianchi I cosmologies with R^n-gravity. *Classical and Quantum Gravity* **23**, 4915 (2006).

[32] Capozziello, S., Nojiri, S., Odintsov, S. & Troisi, A. Cosmological viability of $f(R)$-gravity as an ideal fluid and its compatibility with a matter dominated phase. *Physics Letters B* **639**, 135–143 (2006).

[33] Goheer, N., Larena, J. & Dunsby, P. K. Power-law cosmic expansion in $f(R)$ gravity models. *Physical Review D* **80**, 061301 (2009).

[34] Nojiri, S., Odintsov, S. D. & Sáez-Gómez, D. Cosmological reconstruction of realistic modified $F(R)$ gravities. *Physics Letters B* **681**, 74–80 (2009).

[35] Dunsby, P. K., Elizalde, E., Goswami, R., Odintsov, S. & Saez-Gomez, D. ΛCDM universe in $f(R)$ gravity. *Physical Review D* **82**, 023519 (2010).

[36] Abebe, A. Anti-Newtonian cosmologies in $f(R)$ gravity. *Classical and Quantum Gravity* **31**, 115011 (2014).

[37] Abebe, A. & Elmardi, M. Irrotational-fluid cosmologies in fourth-order gravity. *International Journal of Geometric Methods in Modern Physics* **12**, 1550118 (2015).

[38] Abebe, A., Momeni, D. & Myrzakulov, R. Shear-free anisotropic cosmological models in $f(R)$ gravity. *General Relativity and Gravitation* **48**, 1–17 (2016).

[39] Bean, R., Bernat, D., Pogosian, L., Silvestri, A. & Trodden, M. Dynamics of linear perturbations in $f(R)$ gravity. *Physical Review D* **75**, 064020 (2007).

[40] Song, Y.-S., Hu, W. & Sawicki, I. Large scale structure of $f(R)$ gravity. *Physical Review D* **75**, 044004 (2007).

[41] Abebe, A., Abdelwahab, M., de la Cruz-Dombriz, Á. & Dunsby, P. K. Covariant gauge-invariant perturbations in multifluid $f(R)$ gravity. *Classical and Quantum Gravity* **29**, 135011 (2012).

[42] Carloni, S., Dunsby, P. & Troisi, A. Evolution of density perturbations in $f(R)$ gravity. *Physical Review D* **77**, 024024 (2008).

[43] Ananda, K. N., Carloni, S. & Dunsby, P. K. Structure growth in $f(R)$ theories of gravity with a dust equation of state. *Classical and Quantum Gravity* **26**, 235018 (2009).

[44] Abebe, A., de la Cruz-Dombriz, Á. & Dunsby, P. K. Large scale structure constraints for a class of $f(R)$ theories of gravity. *Physical Review D* **88**, 044050 (2013).

[45] Ananda, K. N., Carloni, S. & Dunsby, P. K. A detailed analysis of structure growth in $f(R)$ theories of gravity. *arXiv preprint arXiv:0809.3673* (2008).

[46] Breaking the cosmological background degeneracy by two-fluid perturbations in $f(R)$ gravity. *Int.J.Mod.Phys. D* **24**, 1550053 (2015).

[47] de La Cruz-Dombriz, A., Dobado, A. & Maroto, A. Evolution of density perturbations in $f(R)$ theories of gravity. *Physical Review D* **77**, 123515 (2008).

[48] Chaplygin, S. On gas jets, Sci. Mem., Moscow Univ. Phys-Math. 21 (1904) pp. 1-127. *Trans. by M. Slud, Brown University* (1944).

[49] Gorini, V., Kamenshchik, A., Moschella, U. & Pasquier, V. The Chaplygin gas as a model for dark energy. In *Proceedings of the MG10 Meeting held at Brazilian Center for Research in Physics (CBPF)*, vol. 20, 26 (World Scientific, 2003).

[50] Bento, M., Bertolami, O. & Sen, A. Generalized Chaplygin gas, accelerated expansion, and dark-energy-matter unification. *Physical Review D* **66**, 043507 (2002).

[51] Bento, M., Bertolami, O. & Sen, A. Letter: Generalized Chaplygin gas model: Dark energy-dark matter unification and cmbr constraints. *General Relativity and Gravitation* **35**, 2063–2069 (2003).

[52] Bilić, N., Tupper, G. B. & Viollier, R. D. Unification of dark matter and dark energy: the inhomogeneous Chaplygin gas. *Physics Letters B* **535**, 17–21 (2002).

[53] Debnath, U., Banerjee, A. & Chakraborty, S. Role of modified Chaplygin gas in accelerated universe. *Classical and Quantum Gravity* **21**, 5609 (2004).

[54] Dev, A., Alcaniz, J. & Jain, D. Cosmological consequences of a Chaplygin gas dark energy. *Physical Review D* **67**, 023515 (2003).

[55] Kamenshchik, A., Moschella, U. & Pasquier, V. An alternative to quintessence. *Physics Letters B* **511**, 265–268 (2001).

[56] Gorini, V., Kamenshchik, A. & Moschella, U. Can the Chaplygin gas be a plausible model for dark energy? *Physical Review D* **67**, 063509 (2003).

[57] Fabris, J. C., Goncurlcalves, S. V. & de Souza, P. E. Letter: density perturbations in a universe dominated by the Chaplygin gas. *General Relativity and Gravitation* **34**, 53–63 (2002).

[58] Bean, R. & Dore, O. Are Chaplygin gases serious contenders for the dark energy? *Physical Review D* **68**, 023515 (2003).

[59] Bouhmadi-López, M., Brilenkov, M., Brilenkov, R., Morais, J. & Zhuk, A. Scalar perturbations in the late universe: viability of the Chaplygin gas models. *Journal of Cosmology and Astroparticle Physics* **2015**, 037 (2015).

[60] Morais, J., Bouhmadi-López, M. & Capozziello, S. Can $f(R)$ gravity contribute to (dark) radiation? *Journal of Cosmology and Astroparticle Physics* **2015**, 041 (2015).

[61] Saadat, H. & Pourhassan, B. Effect of varying bulk viscosity on generalized Chaplygin gas. *International Journal of Theoretical Physics* **53**, 1168–1173 (2014).

[62] Barrow, J. D. Deflationary universes with quadratic lagrangians. *Physics Letters B* **183**, 285–288 (1987).

[63] Makler, M., de Oliveira, S. Q. & Waga, I. Constraints on the generalized Chaplygin gas from supernovae observations. *Physics Letters B* **555**, 1–6 (2003).

[64] Setare, M. Interacting generalized chaplygin gas model in non-flat universe. *The European Physical Journal C* **52**, 689–692 (2007).

[65] Benaoum, H. Accelerated universe from modified Chaplygin gas and tachyonic fluid. *arXiv preprint hep-th/0205140* (2002).

[66] Benaoum, H. Modified Chaplygin gas cosmology. *Advances in High Energy Physics* **2012** (2012).

[67] Pourhassan, B. Viscous modified cosmic Chaplygin gas cosmology. *International Journal of Modern Physics D* **22**, 1350061 (2013).

[68] Kahya, E., Khurshudyan, M., Pourhassan, B., Myrzakulov, R. & Pasqua, A. Higher order corrections of the extended Chaplygin gas cosmology with varying G and Λ. *The European Physical Journal C* **75**, 1–12 (2015).

[69] Kahya, E. & Pourhassan, B. The universe dominated by the extended Chaplygin gas. *Modern Physics Letters A* **30**, 1550070 (2015).

[70] Kahya, E., Pourhassan, B. & Uraz, S. Constructing an inflaton potential by mimicking modified Chaplygin gas. *Physical Review D* **92**, 103511 (2015).

[71] Pourhassan, B. & Kahya, E. FRW cosmology with the extended Chaplygin gas. *Advances in High Energy Physics* **2014** (2014).

[72] Barrow, J. D. & Ottewill, A. C. The stability of general relativistic cosmological theory. *Journal of Physics A: Mathematical and General* **16**, 2757 (1983).